

ON THE RANK OF A SYMMETRIC FORM

KRISTIAN RANESTAD AND FRANK-OLAF SCHREYER

ABSTRACT. We give a lower bound for the degree of a finite apolar subscheme of a symmetric form F , in terms of the degrees of the generators of the annihilator ideal $F^\perp$. In the special case, when F is a monomial $x_0^{d_0} \cdot x_2^{d_2} \cdot \dots \cdot x_n^{d_n}$ with $d_0 \leq d_1 \leq \dots \leq d_{n-1} \leq d_n$ we deduce that the minimal length of an apolar subscheme of F is $(d_0 + 1) \cdot \dots \cdot (d_{n-1} + 1)$, and if $d_0 = \dots = d_n$, then this minimal length coincides with the rank of F .

Let $F \in T = K[x_0, \dots, x_n]$ be a homogeneous form and let $S = K[y_0, \dots, y_n]$ be the ring of commuting differential operators acting on T . The action is called apolarity, and defines S as a natural coordinate ring on the projective space $\mathbf{P}(T_1)$ of 1-dimensional subspaces of T_1 . The annihilator of F is an ideal $F^\perp \subset S$. A finite subscheme $\Gamma \subset \mathbf{P}(T_1)$ is apolar to F if the homogeneous ideal $I_\Gamma \subset S$ is contained in $F^\perp$.

We define the cactus rank $cr(F)$ as

$$cr(F) = \min\{\deg \Gamma \mid \Gamma \subset \mathbf{P}(T_1), \dim \Gamma = 0, I_\Gamma \subset F^\perp\},$$

the smoothable rank $sr(F)$ as

$$sr(F) = \min\{\deg \Gamma \mid \Gamma \subset \mathbf{P}(T_1) \text{ smoothable}, \dim \Gamma = 0, I_\Gamma \subset F^\perp\}$$

and the rank $r(F)$ as

$$r(F) = \min\{\deg \Gamma \mid \Gamma \subset \mathbf{P}(T_1) \text{ smooth}, \dim \Gamma = 0, I_\Gamma \subset F^\perp\}.$$

Clearly $cr(F) \leq sr(F) \leq r(F)$. We shall give lower bounds for these ranks in terms of the generators of the ideal $F^\perp$. The related notion of border rank, $br(F)$, is defined as the minimal k such that $[F]$ lies in the Zariski closure of the set of forms of rank k in $\mathbf{P}(T_{\deg F})$. In general $br(F) \leq sr(F)$, and strict inequality occurs, so our lower bounds for $sr(F)$ does not apply unconditionally to $br(F)$. For applications of these notions of rank to powersum decompositions of symmetric forms and to equations of secant varieties, see [Ranestad, Schreyer 2000], [Landsberg, Teitler 2010] and [Buczynska, Buczyński 2011].

We define the degree of $F^\perp$ to be the length of the quotient algebra $S_F = S/F^\perp$.

Proposition 1. *If the ideal of $F^\perp$ is generated in degree d and $\Gamma \subset \mathbf{P}(T_1)$ is a finite apolar subscheme to F , then*

$$\deg \Gamma \geq \frac{1}{d} \deg F^\perp.$$

Proof. Taking cones, we may assume that $F^\perp$ and I_Γ define subschemes X and Y of pure dimension r and $r + 1$ in $\mathbf{P}^N$. Furthermore $\deg Y = \deg \Gamma$ and $\deg X = \deg F^\perp$. The apolarity condition says that $I_X \supset I_Y$, i.e. that $X \subset Y$ as schemes. Now, take an element g

Date: January 19, 2013.

1991 Mathematics Subject Classification. 13P, 14Xxx.

Key words and phrases. Symmetric tensors.

Both authors were supported by Institut Mittag-Leffler, and thank Jaroslaw Buczyński for clarifying discussions on various notions of rank.

in I_X that does not contain any component of Y . Then the hypersurface $G = \{g = 0\}$ has proper intersection with Y and contains X . Therefore, by Bezout,

$$\deg G \cdot \deg Y \geq \deg X.$$

The proposition follows by taking g of degree d . $\square$

Corollary 1. *If the ideal of $F^\perp$ is generated in degree d , then the cactus rank*

$$cr(F) \geq \frac{1}{d} \deg F^\perp.$$

Corollary 2. *If F is a monomial, $F = x_0^{d_0} \cdot x_1^{d_1} \cdot \dots \cdot x_n^{d_n}$ with $d_0 \leq d_1 \leq \dots \leq d_n$, then the cactus rank and the smoothable rank coincide and equals*

$$cr(F) = sr(F) = (d_0 + 1) \cdot \dots \cdot (d_{n-1} + 1).$$

If furthermore $d_n = d_0 = d$, i.e. $F = (x_0 \cdot x_1 \cdot \dots \cdot x_n)^d$, then $r(F) = cr(F) = sr(F) = (d+1)^n$.

Proof. When $F = x_0^{d_0} \cdot x_1^{d_1} \cdot \dots \cdot x_n^{d_n}$, then $F^\perp$ is the complete intersection generated by the forms

$$y_0^{d_0+1}, y_1^{d_1+1}, \dots, y_n^{d_n+1}.$$

So it is generated in degree $d_n + 1$, while $F^\perp$ has degree

$$(d_0 + 1) \cdot (d_1 + 1) \cdot \dots \cdot (d_n + 1).$$

The formula for the cactus rank follows, since the first n generators define a finite apolar subscheme of degree $(d_0 + 1) \cdot \dots \cdot (d_{n-1} + 1)$. Now, any complete intersection is smoothable, so the smoothable rank equals the cactus rank for F . If $d_0 = d_n = d$, then the forms of degree $d + 1$ in $F^\perp$ has no basepoints so, by Bertini, n general forms in $F^\perp$ of degree $(d + 1)$ define a smooth finite subscheme of degree $(d + 1)^n$ in $\mathbf{P}(T_1)$. $\square$

REFERENCES

- [Buczynska, Buczynski 2011] Buczynska, Weronika, Buczynski, Jaroslaw: Secant varieties to high degree Veronese reembeddings, catalecticant matrices and smoothable Gorenstein schemes, arXiv:1012.3563
- [Landsberg, Teitler 2010] J.M. Landsberg, Zach Teitler, On the ranks and border ranks of symmetric tensors Found Comput Math (2010) 10: 339-366.
- [Ranestad, Schreyer 2000] Ranestad, K., Schreyer, F.-O. Varieties of Sums of Powers. *J. reine angew. Math.* **525**, (2000) 147–181

MATEMATISK INSTITUTT, UNIVERSITETET I OSLO, PO BOX 1053, BLINDERN, NO-0316 OSLO, NORWAY

E-mail address: ranestad@math.uio.no

URL: <http://folk.uio.no/ranestad/>

MATHEMATIK UND INFORMATIK, UNIVERSITÄT DES SAARLANDES, CAMPUS E2 4, D-66123 SAARBRÜCKEN, GERMANY

E-mail address: schreyer@math.uni-sb.de

URL: <http://www.math.uni-sb.de/ag/schreyer/>